

The Value of Liver and Spleen Stiffness for Evaluation of Portal Hypertension in Compensated Cirrhosis

Thomas Reiberger  1-3

Patients with compensated advanced chronic liver disease who develop clinically significant portal hypertension (CSPH) are at high risk for hepatic decompensation and mortality if left untreated. Liver biopsy and hepatic venous pressure gradient (HVPG) measurements are the current gold standard procedures for determining fibrosis severity and diagnosing CSPH, respectively; however, both are invasive, limiting their use in clinical practice and larger trials of novel agents. As such, there is an unmet clinical need for reliable, validated, noninvasive measures to detect CSPH and to further assess portal hypertension (PH) severity. Alterations in the biomechanical properties of the liver or spleen in patients with cirrhosis can be quantified by tissue elastography, which examines the elastic behavior of tissue after a force has been applied. A variety of methods are available, including magnetic resonance elastography, shear-wave elastography, and the most thoroughly investigated measure, vibration-controlled transient elastography. Liver stiffness (LS) and spleen stiffness (SS) measurements offer valuable alternatives to detect and monitor CSPH. Both LS and SS correlate well with HVPG, with thresholds of LS >20-25 kPa and SS >40-45 kPa indicating a high likelihood of CSPH. Because SS is a direct and dynamic surrogate of portal pressure, it has the potential to monitor PH severity and assess PH improvement as a surrogate marker for clinical outcomes. Importantly, SS seems to be superior to LS for monitoring treatment response in clinical trials focusing on reducing PH. (*Hepatology Communications* 2021;0:1-15).

Liver cirrhosis is a consequence of chronic liver disease, with viral hepatitis B (HBV) and hepatitis C (HCV), alcohol-related liver disease (ALD), and nonalcoholic steatohepatitis (NASH) representing the most common etiologies.⁽¹⁾ The Global Burden of Disease study indicates that the prevalence of cirrhosis has almost doubled during the past 3 decades, with more than 122 million cases of cirrhosis documented in 2017.⁽¹⁾ With the use of effective antiviral therapy, HBV and HCV are in

decline yet still represent the leading causes of cirrhosis burden worldwide.⁽¹⁾ However, NASH is increasing in prevalence⁽²⁾ and is soon expected to overtake viral hepatitis to become the leading cause of cirrhosis, hospitalization, and need for liver transplantation.^(1,3) Cirrhosis is clinically classified into two distinct prognostic stages, compensated and decompensated cirrhosis, based on the absence or presence of clinically evident decompensation events that include variceal hemorrhage (VH), hepatic encephalopathy (HE),

Abbreviations: 2D-SWE, two-dimensional shear-wave elastography; ACLD, advanced chronic liver disease; APRI, aminotransferase/platelet ratio index; AST, aspartate aminotransferase; cACLD, compensated advanced chronic liver disease; CSPH, clinically significant portal hypertension; ELF, enhanced liver fibrosis; EV, esophageal varices; FIB-4, fibrosis-4; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HE, hepatic encephalopathy; HVPG, hepatic venous pressure gradient; ICG-r15, indocyanine green 15-minute retention; LS, liver stiffness; MELD, Model for End-Stage Liver Disease; MR, magnetic resonance; NAFLD, nonalcoholic fatty liver disease; NASH, nonalcoholic steatohepatitis; NSBB, nonselective β -blocker; PH, portal hypertension; pSWE, point shear-wave elastography; SS, spleen stiffness; TE, transient elastography; TIPS, transjugular intrahepatic portosystemic shunt; VCTE, vibration-controlled transient elastography; VH, variceal hemorrhage; VNT, varices needing treatment; vWF-Ag, von Willebrand factor antigen.

Received April 25, 2021; accepted October 17, 2021.

Boebring Ingelheim did not have a role in study design, analysis and interpretation of data, writing the manuscript, or the decision to submit the manuscript for publication.

© 2021 The Authors. *Hepatology Communications* published by Wiley Periodicals LLC on behalf of American Association for the Study of Liver Diseases. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

and ascites, the most common first decompensation event.⁽⁴⁾ The compensated stage of cirrhosis is also termed, more comprehensively, compensated advanced chronic liver disease (cACLD) to account for the fact that the transition from advanced fibrosis to cirrhosis is difficult to identify. Importantly, cACLD is further subdivided into two prognostic substages of mild portal hypertension (PH) and clinically significant portal hypertension (CSPH).^(4,5) While patients with CSPH may already show varices at endoscopy, many remain otherwise asymptomatic.^(4,5) Patients may remain in the cACLD stage for decades (median survival >12 years)^(4,5); however, CSPH is the main driver of decompensation.⁽⁶⁾ The decompensated stage is clinically evident by the occurrence of complications related to CSPH and has a much shorter median survival of about 2 years.^(4,5)

Mild PH is a hemodynamic abnormality defined by a hepatic venous pressure gradient (HVPG) of 6–9 mmHg.^(4,7) PH in cirrhosis results from increased vascular resistance caused by structural changes to liver parenchyma (due to collagen deposition, inflammation, nodule formation, and vascular remodeling/occlusion) and increased intrahepatic vascular tone caused by vasoconstriction (driven by decreased vasodilator availability and increased vasoconstrictor production).^(4,8,9)

CSPH is defined by an HVPG of ≥ 10 mmHg and is present in around 60% of patients with cACLD and in all patients with decompensated cirrhosis.^(4,10) CSPH is independently associated with an increased risk of decompensation events, including the development of varices and hepatocellular carcinoma

(HCC).^(4,7,8,11) The risk of developing clinical liver events/decompensation within 2–4 years increases by 11% with each 1 mmHg increase over 10 mmHg in HVPG.^(6,12) Furthermore, reducing HVPG by 20% and/or to <10 mmHg greatly reduces the decompensation risk.⁽¹²⁾ The mortality rate among patients with cirrhosis increases with the development of complications. One-year mortality is 5.4% in patients with compensated cirrhosis but up to 20.2% in patients with decompensated cirrhosis,⁽¹³⁾ while 1-year mortality in patients with compensated cirrhosis without CSPH is 3.4% and increases to 7.3% in patients with CSPH.⁽¹³⁾

Reducing portal pressure in cirrhosis results in better outcomes,⁽¹⁴⁾ and guidelines for the management of cACLD of any etiology focus on prevention of CSPH and decompensation.^(4,7) In patients with compensated cirrhosis, guidelines recommend preventing CSPH by eliminating the underlying cause of liver disease (i.e., etiological therapy) and treating comorbidities that contribute to decompensation or are a consequence of liver disease, such as obesity, diabetes, pulmonary, renal, and cardiovascular disease, to help prevent complications and reduce PH.^(4,7) However, while removing the cause of cirrhosis might improve portal pressure, it is a slow process and patients may continue to be at risk of decompensation.⁽¹⁴⁾

The mainstay of cirrhosis treatment over the last 35 years has been nonselective β -blockers (NSBBs), which act by reducing portal venous inflow.⁽¹⁴⁾ Initially developed to prevent bleeding (and rebleeding) from varices,⁽⁷⁾ a recent randomized, double-blind, placebo-controlled, multicenter trial showed that long-term treatment with

View this article online at wileyonlinelibrary.com.

DOI 10.1002/hep4.1855

Potential conflict of interest: Dr. Reiberger has received grant support from AbbVie, Boehringer Ingelheim, Gilead, MSD, Philips Healthcare, Gore; speaking honoraria from AbbVie, Gilead, Gore, Intercept, Roche, MSD; consulting/advisory board fee from AbbVie, Bayer, Boehringer Ingelheim, Gilead, Intercept, MSD, Siemens; and travel support from AbbVie, Boehringer Ingelheim, Gilead and Roche.

ARTICLE INFORMATION:

From the ¹Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Vienna, Austria; ²Vienna Hepatic Hemodynamic Laboratory, Division of Gastroenterology and Hepatology, Department of Internal Medicine III, Medical University of Vienna, Vienna, Austria; ³Christian-Doppler Laboratory for Portal Hypertension and Liver Fibrosis, Medical University of Vienna, Vienna, Austria.

ADDRESS CORRESPONDENCE AND REPRINT REQUESTS TO:

Thomas Reiberger, M.D.
Division of Gastroenterology and Hepatology, Department of
Medicine III
Medical University of Vienna

Spitalgasse 23
A-1090 Vienna, Austria
E-mail: thomas.reiberger@meduniwien.ac.at
Tel.: +43 1 40400 65890

β -blockers could increase decompensation-free survival in patients with compensated cirrhosis and CSPH, mainly by reducing the incidence of ascites.⁽¹⁵⁾ In a meta-analysis of 15 studies with a total of 1,113 patients with cirrhosis, good hemodynamic response to NSBBs was associated with a lower risk of cirrhosis complications and death or transplantation.⁽¹⁶⁾

For patients who have progressed to decompensated cirrhosis, guidelines recommend treatment to prevent rebleeding of varices and ascites.^(4,7,17,18) Assessment of PH can help to identify those patients at high risk of progression.⁽⁷⁾ There is a need for better therapies to lower portal pressure in cirrhosis, and recent clinical trials of several novel drugs, mainly targeting intrahepatic mechanisms, have shown promise.⁽¹⁴⁾ In phase III clinical trials of cirrhosis, the US Food and Drug Administration currently advocates a composite endpoint based on the occurrence of clinical outcomes (complications of ascites, VH, HE, worsening of Model for End-Stage Liver Disease [MELD] score, liver transplantation, or all-cause death).⁽¹⁹⁾ The European Medicines Agency (EMA) supports the composite endpoint of all-cause death and decompensation events but acknowledges the challenge of defining appropriate endpoints in patients with existing cirrhotic disease.⁽²⁰⁾ For patients with nondecompensated disease, the EMA suggests reversal of cirrhosis (“improvement of liver cirrhosis to noncirrhotic liver disease [1 or more point improvement in fibrosis stage]”) supported by descriptive data on decompensation events, liver transplantation, and death in addition to secondary outcomes derived from noninvasive disease markers, such as biomarkers, imaging data, or determination of liver stiffness (LS), that are the focus of this review.⁽²⁰⁾

Invasive Assessments of Established Prognostic Relevance in cACLD

HVPG

The gold standard for assessing PH is HVPG, which represents the gradient between the pressure in the hepatic sinusoidal capillary network and the free hepatic venous (systemic) pressure. HVPG is measured by retrograde insertion of a balloon-tipped central vein catheter into a main hepatic vein (Fig. 1).^(7,9,21,22)

Guidelines encourage use of HVPG measurements in clinical trials,⁽⁷⁾ and a decrease in HVPG is recommended as the main goal in phase II proof-of-concept studies in PH.⁽¹⁴⁾ If PH-associated endpoints are well defined,⁽⁷⁾ decompensating events are recommended as the main primary endpoint in phase III trials; however, where a low rate of events is expected, HVPG response assessment would be needed as a surrogate marker.^(7,14)

While a decrease in HVPG following treatment with NSBBs (targeting hyperdynamic circulation and splanchnic vasodilation) has been linked to improved outcomes,⁽¹⁶⁾ the association between HVPG change and clinical benefit in developmental therapies for PH may be different and needs to be established.⁽¹⁴⁾ For example, a decrease in HVPG by etiologic therapies for HCV (targeting mainly intrahepatic vascular resistance, the sinusoidal PH) only translates to improved outcomes in patients before PH has developed.⁽²³⁾

LIVER BIOPSY

Liver biopsy is another assessment of cACLD used in the setting of (suspected) cirrhosis, mostly only to differentiate between severe alcoholic hepatitis and decompensated alcoholic cirrhosis⁽²⁴⁾ and to confirm a diagnosis and investigate the possible cause.⁽²⁵⁾ The livers of patients with CSPH in cACLD are modified compared with those with mild PH and develop thick fibrous septa and smaller nodules.⁽⁴⁾ A biopsy may be carried out from a percutaneous or a transjugular route, and while both approaches provide comparable samples, the transjugular route has the advantage that it allows concomitant HVPG measurement.⁽²⁵⁻²⁷⁾ Compared with biopsy, HVPG measurement is better at predicting the likelihood of developing complications of cirrhosis^(4,7); however, both techniques are unsuitable for routine measurements to determine disease progression as both are invasive, and HVPG is not widely available due to high costs and is restricted to highly specialized centers with extensive experience.^(8,11,28) Furthermore, a vasoactive drug may produce rapid, dynamic changes, and repeated measurements in a trial may be required over a number of days. Guidelines, therefore, recommend that research should focus on validation of noninvasive predictors of decompensation for clinical trials.⁽⁷⁾ Some noninvasive techniques have been demonstrated to be of diagnostic value regarding the prediction of risk of CSPH and/or decompensation (Fig. 2).




FIG. 1. Measurement of HVPG. HVPG is measured by retrograde insertion of a balloon-tipped central vein catheter into a main hepatic vein. HVPG represents the difference between the occluded hepatic sinusoidal capillary network (wedged hepatic venous) pressure and the free hepatic venous (systemic) pressure. Abbreviations: FHVP, free hepatic venous pressure; IVCP, inferior vena cava pressure; RAP, right atrium pressure; WHVP, wedged hepatic venous pressure.




FIG. 2. Techniques for identifying CSPH. Multiple screening techniques can be used to identify patients at high risk for progression to decompensated cirrhosis, with various cut-off values to signify the presence of CSPH. Abbreviations: LSM, liver stiffness measurement; SSM, spleen stiffness measurement.

Noninvasive Assessments of Portal Hypertension of Prognostic Relevance in cACLD

A selection of currently used noninvasive assessments of PH are shown in Table 1. Serum levels of degraded extracellular matrix (ECM) products correlate with HVPG and may be regarded as direct biomarkers of PH.⁽²⁹⁾ Platelet counts represent an indirect biomarker of PH, and may be combined with imaging findings (such as spleen diameter) to suggest PH.⁽³⁰⁾ Von Willebrand factor antigen (vWF-Ag) is elevated in patients with liver cirrhosis, correlates with HVPG, is predictive of CSPH, and a value >315% is associated with greater mortality in patients with compensated and decompensated disease.^(31,32) More recently, the vWF-Ag/thrombocyte ratio (VITRO) score, initially derived as a marker of liver fibrosis and cirrhosis in patients with chronic HCV,⁽³³⁾ has been shown to correlate with HVPG and CSPH in patients with cACLD⁽³⁴⁾ and is a predictor of transplant-free mortality and decompensation.⁽³⁵⁾ In patients with liver disease, FibroTest values correlate with HVPG

and the presence and severity of PH, although this association is weaker in patients with cirrhosis.⁽³⁶⁾

Several blood-based indirect fibrosis makers have been investigated in relation to PH. The aspartate aminotransferase (AST)/platelet ratio index (APRI) shows moderate correlation with HVPG, with scores of 0.876 and ≥ 1.09 predicting HVPG ≥ 12 mmHg in two independent studies.^(37,38) The Lok index (derived from AST/alanine aminotransferase ratio, prothrombin and international normalized ratio, and platelet count) and fibrosis-4 (FIB-4) are independently associated with all-cause death in patients with decompensated disease.⁽³⁹⁾ The Lok index has been independently associated with the degree of PH and is predictive of CSPH and esophageal varices (EV).^(40,41) FIB-4 has been associated with the presence of cirrhosis in patients with cACLD.⁽⁴¹⁾ The enhanced liver fibrosis (ELF) score, a combined measurement of serum markers of fibrogenesis and ECM remodeling which was originally developed to detect liver fibrosis, correlates well with HVPG up to values of ≤ 20 mmHg, and an ELF score ≥ 11.1 identifies patients at high risk of CSPH.^(30,42) The indocyanine green 15-minute retention (ICG-r15) test is a quantitative assessment of liver function that can identify CSPH (ICG-r15 $\geq 16.7\%$), rule out the presence of EV (ICG-r15 $> 10\%$),

TABLE 1. NONINVASIVE ASSESSMENTS OF PORTAL HYPERTENSION

Feature	Method	Studies	
Portal venous congestion	LS	Various (see below)	
	SS	Various (see below)	
	Extracellular matrix protein levels (PRO-C3, ELM, C6M)	Leeming et al. ⁽²⁹⁾	
	Platelet count (indirect marker)	Berzigotti et al. ⁽³⁰⁾	
Vasculature	Plasma vWF-Ag	La Mura et al. ⁽³²⁾ Ferlitsch et al. ⁽³¹⁾	
Liver fibrosis	LS	Various (see below)	
	APRI	Verma et al. ⁽³⁷⁾ Kinnake et al. ⁽³⁸⁾	
	Lok index	Lissotti et al. ⁽⁴⁰⁾ Cho et al. ⁽³⁹⁾ Procopet et al. ⁽⁴¹⁾	
	Fib-4	Cho et al. ⁽³⁹⁾ Procopet et al. ⁽⁴¹⁾	
	FibroTest	Thabut et al. ⁽³⁶⁾	
	ELF test	Berzigotti et al. ⁽³⁰⁾ Simbrunner et al. ⁽⁴²⁾	
	Hepatic function	ICG retention	Lisotti et al. ^(40,43)

Abbreviations: C6M, matrix metalloprotease degraded type VI collagen; ELM, elastin; PRO-C3, procollagen type III.

and predict decompensation events (ICG-r15 \geq 23%) in patients with cACLD.^(40,43) Studies have investigated the association between computed tomography and HVPG/CSPH in patients with HCC, HBV-included advanced chronic liver disease (ACLD), and cACLD.⁽⁴⁴⁻⁴⁷⁾ However, a recent review questioned the validity of the findings, highlighting small sample sizes and the inclusion of large proportions of patients with Child-Turcotte-Pugh stage B/C.⁽⁴⁸⁾ Alterations to the biomechanical properties of the liver or spleen in cirrhosis can be measured by elastography techniques, which are based on the elastic behavior of tissue after a force has been applied by ultrasound or magnetic resonance (MR).^(7,8,49)

LIVER STIFFNESS

The main determinant of liver tissue elasticity (also known as LS) is the presence of fibrosis due to collagen deposition, which reflects the main mechanical (“static”) component of hepatic resistance.^(8,50) Initial studies of LS measured with vibration-controlled transient elastography (VCTE) in patients with HCV-related cirrhosis indicated that cut-off values of 13.6 and 17.6 kPa are predictive of HVPG values of \geq 10 and \geq 12 mmHg, respectively.⁽⁵⁰⁾ Subsequent studies have shown reasonable correlation of VCTE with CSPH and HVPG, with a threshold of >20-25 kPa^(51,52); although, it should be noted that in ranges of HVPG values \geq 12 mmHg when complications, such as ascites, develop, this correlation becomes weaker.⁽⁵⁰⁾ Recent guidelines support the noninvasive diagnosis of CSPH by VCTE >20-25 kPa, at least in virus-related cirrhosis.⁽⁷⁾ Similarly, VCTE >20-25 kPa should be used for CSPH risk stratification in cACLD, ideally in combination with platelet count and spleen size.^(4,53) However, for the assessment of PH severity beyond ruling CSPH in or out, current clinical practice guidelines do not suggest that HVPG can be replaced by noninvasive methods and that HVPG remains the only validated tool.⁽⁵³⁾ A meta-analysis of 18 studies concluded that VCTE is able to differentiate between patients with and without CSPH with high accuracy,⁽⁵⁴⁾ whereas another meta-analysis of 11 studies assessing the diagnostic performance of VCTE in assessing CSPH concluded that results of VCTE correlated well with HVPG and proposed cut-off values of 13.6-18 kPa for maximum sensitivity.⁽⁵⁵⁾ In a 2-year study of 41 patients with chronic liver disease, an LS of \geq 21 kPa and an HVPG of \geq 10 mmHg

were equally effective for predicting decompensation events,⁽⁵⁶⁾ and a study of 78 patients with chronic liver disease demonstrated that LS, measured by point shear-wave elastography (pSWE), could identify an HVPG of \geq 10 mmHg and \geq 12 mmHg with high diagnostic performance.⁽⁵⁷⁾ The Anticipate study including 518 patients with cACLD concluded that VCTE and LS to spleen/platelet score (LSPS) reliably identifies patients at high risk of CSPH, with LSPS providing the highest level of discrimination.⁽⁵⁸⁾ In a study of patients with cACLD secondary to nonalcoholic fatty liver disease (NAFLD), an LS of 21.8 kPa identified esophageal varices with high accuracy and has been proposed as a threshold that may obviate the need for a diagnostic esophagogastroduodenoscopy⁽⁵⁹⁾ in line with existing recommendations (<20 kPa) for patients with cACLD of viral origin.⁽⁷⁾ A summary of the heterogeneity of cut-off values for LS that have been proposed to identify HVPG \geq 10 and \geq 12 mmHg and other clinically relevant outcomes related to PH, such as presence of EVs or varices needing treatment (VNT), are listed in Table 2. Ultimately, it is evident that a wide range of published cut-off values for ruling in and ruling out cACLD and CSPH result from heterogeneous patient characteristics with variable pretest and posttest probabilities of the condition of interest. Thus, the Baveno VI consensus guidelines recommend the following use of “easy-to-remember” LS measurement cut-off values for daily clinical use: (i) at LS <10 kPa, rule out cACLD if no other clinical signs, (ii) an LS >15 kPa is highly suggestive of cACLD, and (iii) at LS \geq 20-25 kPa, rule in CSPH in patients with virus-related cACLD.⁽⁷⁾ Importantly, these cut-off values have been mostly validated in patients with hepatitis C,⁽⁶⁰⁾ and although supporting evidence has been created in patients with cholestatic liver disease,⁽⁶¹⁾ historically these cut-off values did not perform as well in patients with NASH due to the close association with obesity.⁽⁶²⁾ High body mass index (BMI), particularly \geq 30 kg/m², is independently associated with higher LS measurements, affecting the interpretation of LS cut-off values when using the standard probe.⁽⁶³⁾ However, development of the XL probe has enabled the use of standard cut-off values in patients with obesity.⁽⁶⁴⁾

SPLEEN STIFFNESS

Since the splenic blood flow drains via the splenic vein into the portal venous system, spleen stiffness (SS) not only reflects static hepatic resistance due to liver

TABLE 2. LS CUT-OFF VALUES

Study	Etiology	Modality	HVPG ≥ 10 mmHg	HVPG ≥ 12 mmHg	Other
Vizzutti et al. ⁽⁵⁰⁾	HCV n = 61	VCTE	13.6 kPa	17.6 kPa	EV: 17.6 kPa
Bureau et al. ⁽¹⁰⁵⁾	CLD n = 150	VCTE	21 kPa		EV: 21.1 kPa
Lemoine et al. ⁽¹⁰⁶⁾	HCV n = 44	VCTE	20.5 kPa		
	ALD n = 48	VCTE	34.9 kPa		
Sanchez-Conde et al. ⁽¹⁰⁷⁾	HIV-HCV n = 100	VCTE	14.0 kPa	23.0 kPa	
Robic et al. ⁽⁵⁶⁾	CLD n = 100	VCTE			Low-risk CD: ≤ 21.1 kPa High-risk CD: > 21.1 kPa
Colecchia et al. ⁽²⁸⁾	HCV n = 100	VCTE	24.2 kPa	25.0 kPa	EV: 25.0 kPa
Reiberger et al. ⁽¹⁰⁸⁾	HCV n = 390	VCTE	18.0 kPa	20.0 kPa	Mild PH (≥ 6 mmHg): 8.0 kPa
	ALD n = 227	VCTE	19.0 kPa	23.0 kPa	Mild PH (≥ 6 mmHg): 10.0 kPa
Llop et al. ⁽¹⁰⁹⁾	HCC n = 97	VCTE	21 kPa		
Hong et al. ⁽¹¹⁰⁾	cACLD n = 59	VCTE	21.95 kPa	24.25 kPa	
Sharma et al. ⁽¹¹¹⁾	cACLD n = 174	VCTE			EV: 27.3 kPa
Augustin et al. ⁽⁵²⁾	CLD n = 250	VCTE	25 kPa		
Salzl et al. ⁽⁸³⁾	cACLD n = 88	VCTE	16.8 kPa		EV: 27.9 kPa
Attia et al. ⁽⁵⁷⁾	CLD n = 78	pSWE	2.58 m/second		EV: 2.74 m/second
		pSWE	2.17 m/second		
Elkrief et al. ⁽⁸⁴⁾	cACLD n = 79	VCTE	65.3 kPa		
		SWE	24.5 kPa		
Kim et al. ⁽¹¹²⁾	cACLD n = 115	SWE	15.2 kPa	21.6 kPa	
Procopet et al. ⁽⁸⁶⁾	CLD n = 202	VCTE	13.6 kPa		
		SWE	15.4 kPa		
Schwabl et al. ⁽¹¹³⁾	CLD n = 226	VCTE	16.1 kPa		Cirrhosis: 14.5 kPa
Jansen et al. ⁽⁹⁰⁾	ACLD n = 158	SWE	16.0 kPa		Rule out CSPH: LSM < 16 kPa and SS < 26.6 kPa Rule in CSPH: LSM ≥ 16 kPa or SS ≥ 26.6 kPa
Jansen et al. ⁽¹¹⁴⁾	cACLD n = 158	SWE	24.6 kPa	28.5 kPa	Rule out CSPH: LSM 16.0 kPa Rule in CSPH: LSM > 29.5 kPa or SSM > 27.9 kPa
Wong et al. ⁽⁶⁴⁾	NAFLD n = 548	VCTE			Rule out cACLD: < 10 kPa Rule in cACLD: ≥ 15 kPa
Galizzi et al. ⁽⁵⁹⁾	cACLD-NAFLD n = 21	VCTE			Any EV: 21.8 kPa VNT: 21.8 kPa

TABLE 2. Continued

Study	Etiology	Modality	HVPG ≥ 10 mmHg	HVPG ≥ 12 mmHg	Other
Souhami et al. ⁽¹¹⁵⁾	HCC n = 140	VCTE	21.0 kPa		
Stefanescu et al. ⁽⁷¹⁾	cACLD n = 127	SWE	11.3 kPa		
Thiele et al. ⁽⁸⁵⁾	ACLD n = 328	2D-SWE			Rule out CSPH: <14 kPa
Trebicka et al. ⁽⁸⁷⁾	ACLD n = 1,827	2D-SWE			High risk of decompensation or death: ≥ 20 kPa and MELD ≥ 10

Abbreviations: ALD, alcohol-related liver disease; CD, clinical decomposition; CLD, chronic liver disease; HIV, human immunodeficiency virus.

fibrosis (that is captured by LS) but may also capture dynamic (pre)sinusoidal vasoconstriction and congestion of the portal blood inflow (a potential surrogate of PH-associated splanchnic hypoperfusion) and PH-induced splenic fibrosis.⁽⁶⁵⁻⁶⁸⁾ SS is impacted by an increase in PH, which leads to splenic congestion, but also induces architectural changes in the splenic arteries and veins and fibrosis.⁽⁶⁶⁾ Thus, SS measurement is a valuable surrogate of PH (and initially an SS value of 56 kPa was reported to be associated with CSPH), and SS correlates well with HVPG irrespective of the etiology.^(28,69,70) However, as evidence has accumulated, it appears that lower thresholds of ≤ 41 - 46 kPa are able to rule out the presence of CSPH and high-risk varices.^(28,71-73) In a meta-analysis of nine studies, SS measured by ultrasound-based elastography showed good correlation with HVPG, detecting CSPH with a sensitivity and specificity of 0.88 and 0.92, respectively, and severe PH with a sensitivity and specificity of 0.92 and 0.79, respectively.⁽⁷⁰⁾ A study of 100 patients with HCV also demonstrated that SS, measured by transient elastography (TE), correlated strongly with the whole range of HVPG values >5 mmHg, suggesting that the increase in SS is closely tied with the progression of PH from early to late stages of cirrhosis.⁽²⁸⁾ SS has been combined with MELD score to provide an accurate predictor of decompensation in patients with HCV and is at least as accurate as HVPG.⁽⁶⁹⁾ In patients with HCV-related ACLD, SS is a direct surrogate of persisting PH following HCV therapy with direct-acting antivirals.⁽¹¹⁾ Studies of healthy volunteers and patients with various etiologies of chronic liver disease indicate that SS, as measured by MR elastography, correlates with platelet count, spleen volume, and splenomegaly and

that an SS of ≥ 10.5 kPa is associated with esophageal varices.⁽⁷⁴⁾ SS as determined by pSWE has also been shown to be predictive of the development of esophageal varices,⁽⁷⁵⁾ and in a study of 78 patients with chronic liver disease it was able to identify an HVPG of ≥ 10 mmHg and ≥ 12 mmHg with high diagnostic performance.⁽⁵⁷⁾ The different cut-off values of SS that have been proposed to predict HVPG ≥ 10 and ≥ 12 mmHg and other clinically relevant outcomes related to PH, such as the detection of high-risk EV, are summarized in Table 3.

LIVER STIFFNESS VERSUS SPLEEN STIFFNESS

Data show that SS may be a superior marker of PH to LS in patients with cirrhosis with viral etiologies, and an increasing body of evidence suggests this may also be the case for other etiologies.^(11,28,69,76,77) For example, in a meta-analysis of 16 studies of chronic liver disease, SS detected esophageal varices (one of the most serious consequences of PH⁽⁷⁸⁾ that is independently predicted by HVPG⁽¹⁰⁾) with a superior sensitivity and specificity (0.88 and 0.78, respectively) to LS (0.83 and 0.66, respectively).⁽⁷⁷⁾ Furthermore, in studies of virus-related compensated cirrhosis, SS was better than LS and similar to HVPG in predicting a first clinical decompensation event.^(28,69) A recent study also showed that compared with LS, SS measurement was a direct surrogate of PH in patients after cure from HCV and may be useful for monitoring response and stratifying risk following therapy.⁽¹¹⁾ This may be explained by the fact that SS measurements are less influenced by liver necroinflammation than LS measurements.⁽¹¹⁾ In addition, SS might be

TABLE 3. SS CUT-OFF VALUES

Study	Etiology	Modality	HVPG \geq 10 mmHg	HVPG \geq 12 mmHg	Other
Talwalkar et al. ⁽⁷⁴⁾	CLD n = 38	MRE			EV: 10.5 kPa
Colecchia et al. ⁽²⁸⁾	HCV n = 100	VCTE	52.8 kPa	55.0 kPa	EV: 55.0 kPa
Sharma et al. ⁽¹¹¹⁾	cACLD n = 174	VCTE			EV: 40.8 kPa
Colecchia et al. ⁽⁶⁹⁾	HCV n = 92	VCTE			Low risk of CD: <54.0 kPa High risk of CD: >54.0 kPa
Elkrief et al. ⁽⁸⁴⁾	cACLD n = 79	VCTE SWE	56.3 kPa 34.7 kPa		
Kim et al. ⁽¹¹⁶⁾	cACLD n = 125	pSWE			EV: 3.16 m/second
Jansen et al. ⁽⁹⁰⁾	ACLD n = 158	SWE			CSPH: ≥ 26.6 kPa and LS < 1.6 kPa
Jansen et al. ⁽¹¹⁴⁾	cACLD n = 158	SWE	26.3 kPa	28.5 kPa	
Colecchia et al. ⁽⁷²⁾	cACLD n = 498	VCTE			Rule out VNT: ≤ 46 kPa
Fierbinteanu-Braticevici et al. ⁽⁷⁵⁾	cACLD n = 135	pSWE			Any EV: 3m/second VNT: 3.5 m/second
Marasco et al. ⁽¹¹⁷⁾	CLD after HCC n = 157	VCTE			Late HCC recurrence: >70 kPa
Marasco et al. ⁽⁸¹⁾	ACLD with HRV n = 20	VCTE			No NSBB: 61.5 kPa With NSBB: 35.8 kPa
Stefanescu et al. ⁽¹¹⁸⁾	ACLD n = 260	Spleen-TE	34.16 kPa	44.95 kPa	HRV: 41.3 kPa
Cho et al. ⁽¹¹⁹⁾	cACLD n = 270	SWE			Rule out HRV: ≤ 27.3 kPa
Wang et al. ⁽¹⁰⁴⁾	HBV, cACLD n = 341	VCTE			Rule out HRV: ≤ 46 kPa
Dajti et al. ⁽⁷³⁾	ACLD				Low-risk VNT: ≤ 46 kPa
Dajti et al. ⁽¹²⁰⁾	HCV-related ACLD n = 140	VCTE			High-risk HCC: >42 kPa

Abbreviations: CD, clinical decomposition; HRV, high-risk varices; MRE, magnetic resonance elastography.

a more dynamic marker, reflecting “acute” changes in HVPG that are not detected by LS (e.g., a decrease in portal pressure occurring immediately after and within 30 minutes of transjugular intrahepatic portosystemic shunt [TIPS]⁽⁷⁹⁾) or changes in HVPG that occur mainly due to pharmacologic inhibition of hyperdynamic circulation/splanchnic blood flow.⁽⁸⁰⁾ For example, in a single-cohort, proof-of-concept study in patients with CSPH in cirrhosis and high-risk varices, a change in SS predicted a hemodynamic response ($\geq 20\%$ decrease in HVPG or an HVPG

< 12 mmHg) to the NSBB carvedilol whereas LS and HVPG did not.⁽⁷⁶⁾ Another study investigated changes in SS measured by TE in 20 patients with cirrhosis and high-risk varices undergoing sequential HVPG measurement before and during NSBB therapy.⁽⁸¹⁾ The authors found that SS significantly correlated with changes in HVPG following NSBB therapy ($r = 0.784$; $P < 0.0001$) while LS measured by TE did not ($r = 0.107$, $P = 0.655$).⁽⁸¹⁾ Importantly, a $\geq 10\%$ decrease in SS was an excellent predictor of HVPG response to NSBBs (area under the receiver

operator characteristic curve, 0.973; 95% confidence interval, 0.912-1.000) and was associated with a sensitivity of 100% and specificity of 60%.⁽⁸¹⁾ However, due to the limited sample sizes and uncontrolled designs, additional validation studies of SS are required.⁽⁸²⁾

American Association for the Study of Liver Diseases practice guidelines (2016) state that newer sonoelastography techniques, such as pSWE, can facilitate SS measurements and compared with TE have similar or greater accuracy in predicting CSPH.^(4,83,84) Two-dimensional shear-wave elastography (2D-SWE) can be applied to LS and SS, with an LS cut-off value of 14 kPa able to rule out the presence of CSPH in patients with cACLD.⁽⁸⁵⁾ LS by 2D-SWE is superior to SS by 2D-SWE, correlates strongly with VCTE, and can accurately diagnose CSPH if suitable reliability criteria are fulfilled.⁽⁸⁶⁾ A large international study used a 2D-SWE LS cut-off value of ≥ 20 kPa combined with a MELD score of ≥ 10 (the M10LS20 algorithm) to stratify the risk of decompensation and mortality in patients with compensated or decompensated cirrhosis.⁽⁸⁶⁾ This algorithm was validated by both TE and pSWE measurements and can identify patients requiring intensified care and early treatment.⁽⁸⁶⁾ However, although these techniques show promise, the heterogeneity of cut-off values observed and the limited number of studies mean their use cannot yet be routinely recommended in clinical practice guidelines.⁽⁵³⁾

LS and SS can be used in combination to identify high-risk patients and rule out the presence of CSPH. A randomized, controlled trial showed the potential noninferiority of combined LS and SS measurements compared with universal endoscopic screening for detecting clinically significant varices in patients with cirrhosis.⁽⁸⁸⁾ A prospective follow-up study showed that patients undergoing this screening technique were at low risk for incident VH and that the use of this technique reduced the need for endoscopic screening by up to 50%.⁽⁸⁹⁾ Another study developed an algorithm to identify patients with CSPH by using sequential LS and SS measurements by SWE, with cut-off values of LS ≥ 16.0 kPa or LS < 16.0 kPa and SS ≥ 26.6 kPa showing a sensitivity of 98.6%.⁽⁹⁰⁾ As LS is the most validated noninvasive test for PH, clinical practice guidelines only currently recommend SS as an additional measure that can improve risk stratification for high-risk varices and CSPH.⁽⁵³⁾

DETERMINANTS AND CONFOUNDERS OF TISSUE STIFFNESS

In addition to accuracy, reliability and repeatability of measurements are important factors in assessing the performance of sonoelastography techniques. The ability to achieve consistent and reproducible measurements depends on several critical factors, and multiple confounders affect both LS and SS measurements and can lead to false-positive diagnoses of fibrosis and CSPH.

First, the ability to achieve consistent and reproducible results can depend on the experience of the operator, with a greater impact on SS than LS.⁽⁹¹⁾ Although operator experience is more relevant with pSWE and 2D-SWE than with VCTE, inexperience also decreases the reliability of LS measured by VCTE.⁽⁹²⁾

Second, obesity (mostly due to a higher skin-to-liver capsule distance) impacts LS results. In a 5-year study of 13,369 VCTE-based LS examinations in 7,261 patients, almost one in five measurements were unreliable due to the impact of body weight and increased waist circumference, making its application limited in patients affected by obesity.⁽⁹²⁾ Importantly, a VCTE probe specifically designed for patients with obesity (i.e., the XL probe) has been developed,⁽⁹¹⁾ allowing for LS measurements at a greater depth. This XL probe yields accurate VCTE measurements in patients with obesity that are similar to measurements obtained with a standard probe in patients without obesity.⁽⁶⁴⁾ This allows the same VCTE cut-off values to be used for cACLD diagnosis in patients with higher BMI.⁽⁶⁴⁾

Third, venous congestions of the liver (i.e., due to heart failure) also increase LS up to values that clearly fall in the "cirrhotic" range, thereby precluding fibrosis staging in patients with heart failure (such as in Fontan-associated liver failure).^(93,94)

Fourth, mechanical cholestasis increases LS,⁽⁹⁵⁾ with bile duct obstruction leading to increased pressure, edema, swelling, and inflammation that could lead to incorrect patient classification.⁽⁹⁶⁾

Fifth, postprandial increases in portal venous perfusions impact LS results,^(97,98) with 10%-16% of patients with cirrhosis misclassified with CSPH after the consumption of a moderate- or high-calorie meal.⁽⁹⁹⁾ SS was also significantly ($P < 0.001$) increased

from baseline after food intake by 17%–19%.⁽⁹⁹⁾ These elevated measurements can persist for 3 hours after a meal, leading to recommendations of at least a >3-hour fasting period before LS or SS measurements.⁽⁹⁹⁾

Sixth, local and systemic inflammation increases LS, likely due to inflammatory cell infiltrations and hepatocellular swelling.^(100,101) Inflammation-dependent LS cut-off values have been developed based on increased AST levels to improve fibrosis staging and could be used to identify patients at high risk irrespective of inflammation.⁽¹⁰¹⁾ However, the effect of inflammation on SS is less clear, with studies suggesting SS measurements may be less biased by inflammation than LS,⁽¹⁰²⁾ but aminotransferase levels can still be a significant confounding factor.⁽¹⁰³⁾ One study looked at the effect of inflammation on LS and SS in patients receiving TIPS for PH.⁽⁷⁹⁾ This study found that, in patients with higher levels of inflammatory markers, LS increased after TIPS despite the decrease in portal pressure observed; this effect was not seen in SS.⁽⁷⁹⁾

SS measurements using the conventional TE probe are difficult, with a higher failure rate for SS than LS.⁽⁹¹⁾ In a prospective study of 341 patients with HBV, reliable SS could not be obtained in 4.1% of patients (14/341) due to factors relating to obesity, ascites, or invalid SS measurements.⁽¹⁰⁴⁾ Similarly, in patients with compensated disease (cACLD without ascites), SS failed in 5.2% of patients (26/498).⁽⁷²⁾ Specific training has been shown to improve the repeatability of SS measurements.⁽⁹¹⁾ More recently, a novel spleen-dedicated stiffness measurement TE probe has been developed that has further improved the screening for high-risk esophageal varices.⁽⁷¹⁾

To compensate for the potential pitfall of false-positive results, it is recommended to repeat VCTE LS measurements on at least 2 different days under fasting conditions and not use LS for fibrosis staging in patients presenting with venous congestion, mechanical cholestasis, pronounced inflammation, and high levels of transaminases.⁽⁷⁾

Conclusions

Patients with compensated cirrhosis (i.e., cACLD) who develop CSPH are at high risk for progression to decompensated cirrhosis and have increased mortality if left untreated. With limited treatment options

available, there is an unmet clinical need to develop effective therapies to lower portal pressure in cirrhosis. Clinical trials for novel therapies require validated outcome measures that are easily repeatable. Current procedures for determining PH severity in cACLD, such as HVPG and biopsy, are invasive, expensive, and restricted to specialist centers, which limit their repeatability and use both in clinical practice and in larger trials. Noninvasive measurements of LS and SS correlate well with HVPG and offer valuable alternatives to monitor CSPH. Because SS is a direct and dynamic surrogate of PH that has the potential to assess an improvement in PH as a surrogate marker for clinical outcomes, SS may be a superior tool to LS for monitoring therapy response in clinical trials.

Acknowledgment: I thank Juan Gonzalez Abraldes for useful discussions during the initial stages of manuscript development. Medical writing support was provided by Tom Priddle, D.Phil., and Susie Eaton, M.Bio., of OPEN Health Communications (London, United Kingdom), funded by Boehringer Ingelheim.

REFERENCES

- 1) GBD 2017 Cirrhosis Collaborators. The global, regional, and national burden of cirrhosis by cause in 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Gastroenterol Hepatol* 2020;5: 245–266.
- 2) Estes C, Anstee QM, Arias-Loste MT, Bantel H, Bellentani S, Caballeria J, et al. Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016–2030. *J Hepatol* 2018;69: 896–904.
- 3) Axley P, Ahmed Z, Arora S, Haas A, Kuo Y-F, Kamath PS, et al. NASH is the most rapidly growing etiology for acute-on-chronic liver failure-related hospitalization and disease burden in the United States: a population-based study. *Liver Transpl* 2019;25:695–705.
- 4) Garcia-Tsao G, Abraldes JG, Berzigotti A, Bosch J. Portal hypertensive bleeding in cirrhosis: risk stratification, diagnosis, and management: 2016 practice guidance by the American Association for the Study of Liver Diseases. *Hepatology* 2017;65: 310–335. Erratum in: *Hepatology* 2017;66:304.
- 5) D'Amico G, Garcia-Tsao G, Pagliaro L. Natural history and prognostic indicators of survival in cirrhosis: a systematic review of 118 studies. *J Hepatol* 2006;44:217–231.
- 6) Ripoll C, Groszmann R, Garcia-Tsao G, Grace N, Burroughs A, Planas R, et al.; Portal Hypertension Collaborative Group. Hepatic venous pressure gradient predicts clinical decompensation in patients with compensated cirrhosis. *Gastroenterology* 2007;133:481–488.
- 7) de Franchis R; Baveno VI Faculty. Expanding consensus in portal hypertension: report of the Baveno VI Consensus Workshop: stratifying risk and individualizing care for portal hypertension. *J Hepatol* 2015;63:743–752.

- 8) Roccarina D, Rosselli M, Genesca J, Tsochatzis EA. Elastography methods for the non-invasive assessment of portal hypertension. *Expert Rev Gastroenterol Hepatol* 2018;12:155-164.
- 9) Simonetto DA, Liu M, Kamath PS. Portal hypertension and related complications: diagnosis and management. *Mayo Clin Proc* 2019;94:714-726.
- 10) Groszmann RJ, Garcia-Tsao G, Bosch J, Grace ND, Burroughs AK, Planas R, et al.; Portal Hypertension Collaborative Group. Beta-blockers to prevent gastroesophageal varices in patients with cirrhosis. *N Engl J Med* 2005;353:2254-2261.
- 11) Ravaioli F, Colecchia A, Dajti E, Marasco G, Alemanni LV, Tamè M, et al. Spleen stiffness mirrors changes in portal hypertension after successful interferon-free therapy in chronic-hepatitis C virus patients. *World J Hepatol* 2018;10:731-742.
- 12) Sanyal AJ, Harrison SA, Ratziu V, Abdelmalek MF, Diehl AM, Caldwell S, et al. The natural history of advanced fibrosis due to nonalcoholic steatohepatitis: data from the simtuzumab trials. *Hepatology* 2019;70:1913-1927.
- 13) Zipprich A, Garcia-Tsao G, Rogowski S, Fleig WE, Seufferlein T, Dollinger MM. Prognostic indicators of survival in patients with compensated and decompensated cirrhosis. *Liver Int* 2012;32:1407-1414.
- 14) Abraldes JG, Trebicka J, Chalasani N, D'Amico G, Rockey DC, Shah VH, et al. Prioritization of therapeutic targets and trial design in cirrhotic portal hypertension. *Hepatology* 2019;69:1287-1299.
- 15) Villanueva C, Albillos A, Genesca J, Garcia-Pagan JC, Calleja JL, Aracil C, et al. β blockers to prevent decompensation of cirrhosis in patients with clinically significant portal hypertension (PREDESCI): a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet* 2019;393:1597-1608. Erratum in: *Lancet* 2019;393:2492.
- 16) Turco L, Villanueva C, La Mura V, García-Pagán JC, Reiberger T, Genesca J, et al. Lowering portal pressure improves outcomes of patients with cirrhosis, with or without ascites: a meta-analysis. *Clin Gastroenterol Hepatol* 2020;18:313-327.e316.
- 17) European Association for the Study of the Liver. EASL Clinical Practice Guidelines for the management of patients with decompensated cirrhosis. *J Hepatol* 2018;69:406-460.
- 18) Ge PS, Runyon BA. Treatment of patients with cirrhosis. *N Engl J Med* 2016;375:767-777.
- 19) U.S. Department of Health and Human Services; Food and Drug Administration; Center for Drug Evaluation and Research (CDER). Nonalcoholic steatohepatitis with compensated cirrhosis: developing drugs for treatment. Guidance for industry. <https://www.fda.gov/media/127738/download>. Published June 2019. Accessed October 2020.
- 20) European Medicines Agency Committee for Medicinal Products for Human Use (CHMP). Reflection paper on regulatory requirements for the development of medicinal products for chronic non-infectious liver diseases (PBC, PSC, NASH). https://www.ema.europa.eu/en/documents/scientific-guideline/reflection-paper-regulatory-requirements-development-medicinal-products-chronic-non-infectious-liver_en.pdf. Published November 15, 2018. Accessed October 2020.
- 21) Paton A, Reynolds TB, Sherlock S. Assessment of portal venous hypertension by catheterisation of hepatic vein. *Lancet* 1953;1:918-921.
- 22) Suk KT. Hepatic venous pressure gradient: clinical use in chronic liver disease. *Clin Mol Hepatol* 2014;20:6-14.
- 23) Libânio D, Marinho RT. Impact of hepatitis C oral therapy in portal hypertension. *World J Gastroenterol* 2017;23:4669-4674.
- 24) European Association for the Study of Liver. EASL clinical practical guidelines: management of alcoholic liver disease. *J Hepatol* 2012;57:399-420.
- 25) Procopet B, Berzigotti A. Diagnosis of cirrhosis and portal hypertension: imaging, non-invasive markers of fibrosis and liver biopsy. *Gastroenterol Rep (Oxf)* 2017;5:79-89.
- 26) Kalambokis G, Manousou P, Vibhakorn S, Marelli L, Cholongitas E, Senzolo M, et al. Transjugular liver biopsy—indications, adequacy, quality of specimens, and complications—a systematic review. *J Hepatol* 2007;47:284-294.
- 27) Reiberger T, Schwabl P, Trauner M, Peck-Radosavljevic M, Mandorfer M. Measurement of the hepatic venous pressure gradient and transjugular liver biopsy. *J Vis Exp* 2020;160:e58819.
- 28) Colecchia A, Montrone L, Scaiola E, Bacchi-Reggiani ML, Colli A, Casazza G, et al. Measurement of spleen stiffness to evaluate portal hypertension and the presence of esophageal varices in patients with HCV-related cirrhosis. *Gastroenterology* 2012;143:646-654.
- 29) Leeming DJ, Karsdal MA, Byrjalsen I, Bendtsen F, Trebicka J, Nielsen MJ, et al. Novel serological neo-epitope markers of extracellular matrix proteins for the detection of portal hypertension. *Aliment Pharmacol Ther* 2013;38:1086-1096.
- 30) Berzigotti A, Seijo S, Arena U, Abraldes JG, Vizzutti F, García-Pagán JC, et al. Elastography, spleen size, and platelet count identify portal hypertension in patients with compensated cirrhosis. *Gastroenterology* 2013;144:102-111.e101.
- 31) Ferlitsch M, Reiberger T, Hoke M, Salzl P, Schwengerer B, Ulbrich G, et al. von Willebrand factor as new noninvasive predictor of portal hypertension, decompensation and mortality in patients with liver cirrhosis. *Hepatology* 2012;56:1439-1447.
- 32) La Mura V, Reverter JC, Flores-Arroyo A, Raffa S, Reverter E, Seijo S, et al. Von Willebrand factor levels predict clinical outcome in patients with cirrhosis and portal hypertension. *Gut* 2011;60:1133-1138.
- 33) Maieron A, Salzl P, Peck-Radosavljevic M, Trauner M, Hametner S, Schöfl R, et al. Von Willebrand Factor as a new marker for non-invasive assessment of liver fibrosis and cirrhosis in patients with chronic hepatitis C. *Aliment Pharmacol Ther* 2014;39:331-338.
- 34) Hametner S, Ferlitsch A, Ferlitsch M, Etschmaier A, Schöfl R, Ziachehabi A, et al. The VITRO score (von Willebrand factor antigen/thrombocyte ratio) as a new marker for clinically significant portal hypertension in comparison to other non-invasive parameters of fibrosis including ELF test. *PLoS One* 2016;11:e0149230.
- 35) Schwarzer R, Reiberger T, Mandorfer M, Kivaranovic D, Hametner S, Hametner S, et al. The von Willebrand factor antigen to platelet ratio (VITRO) score predicts hepatic decompensation and mortality in cirrhosis. *J Gastroenterol* 2020;55:533-542.
- 36) Thabut D, Imbert-bismut F, Cazals-hatem D, Messous D, Muntenau M, Valla DC, et al. Relationship between the Fibrotest and portal hypertension in patients with liver disease. *Aliment Pharmacol Ther* 2007;26:359-368.
- 37) Verma V, Sarin SK, Sharma P, Kumar A. Correlation of aspartate aminotransferase/platelet ratio index with hepatic venous pressure gradient in cirrhosis. *United European Gastroenterol J* 2014;2:226-231.
- 38) Kirnake V, Arora A, Sharma P, Goyal M, Chawlani R, Toshniwal J, et al. Non-invasive aspartate aminotransferase to platelet ratio index correlates well with invasive hepatic venous pressure gradient in cirrhosis. *Indian J Gastroenterol* 2018;37:335-341.
- 39) Cho EJ, Kim MY, Lee J-H, Lee IY, Lim YL, Choi DH, et al. Diagnostic and prognostic values of noninvasive predictors of portal hypertension in patients with alcoholic cirrhosis. *PLoS One* 2015;10:e0133935.
- 40) Lisotti A, Azzaroli F, Buonfiglioli F, Montagnani M, Cecinato P, Turco L, et al. Indocyanine green retention test as a noninvasive

- marker of portal hypertension and esophageal varices in compensated liver cirrhosis. *Hepatology* 2014;59:643-650.
- 41) Procopet B, Cristea VM, Robic MA, Grigorescu M, Agachi PS, Metivier S, et al. Serum tests, liver stiffness and artificial neural networks for diagnosing cirrhosis and portal hypertension. *Dig Liver Dis* 2015;47:411-416.
 - 42) Simbrunner B, Marculescu R, Scheiner B, Schwabl P, Bucsecs T, Stadlmann A, et al. Non-invasive detection of portal hypertension by enhanced liver fibrosis score in patients with different aetiologies of advanced chronic liver disease. *Liver Int* 2020;40:1713-1724.
 - 43) Lisotti A, Azzaroli F, Cucchetti A, Buonfiglioli F, Cecinato P, Calvanese C, et al. Relationship between indocyanine green retention test, decompensation and survival in patients with Child-Pugh A cirrhosis and portal hypertension. *Liver Int* 2016;36:1313-1321.
 - 44) Iranmanesh P, Vazquez O, Terraz S, Majno P, Spahr L, Poncet A, et al. Accurate computed tomography-based portal pressure assessment in patients with hepatocellular carcinoma. *J Hepatol* 2014;60:969-974.
 - 45) Qi X, An W, Liu F, Qi R, Wang L, Liu Y, et al. Virtual hepatic venous pressure gradient with CT angiography (CHESS 1601): a prospective multicenter study for the noninvasive diagnosis of portal hypertension. *Radiology* 2019;290:370-377.
 - 46) Liu F, Ning Z, Liu Y, Liu D, Tian J, Luo H, et al. Development and validation of a radiomics signature for clinically significant portal hypertension in cirrhosis (CHESS1701): a prospective multicenter study. *EBioMedicine* 2018;36:151-158.
 - 47) Sartoris R, Rautou P-E, Elkrief L, Pollorsi G, Durand F, Valla D, et al. Quantification of liver surface nodularity at CT: utility for detection of portal hypertension. *Radiology* 2018;289:698-707.
 - 48) Mandorfer M, Hernández-Gea V, García-Pagán JC, Reiberger T. Noninvasive diagnostics for portal hypertension: a comprehensive review. *Semin Liver Dis* 2020;40:240-255.
 - 49) Ravaoli F, Montagnani M, Lisotti A, Festi D, Mazzella G, Azzaroli F. Noninvasive assessment of portal hypertension in advanced chronic liver disease: an update. *Gastroenterol Res Pract* 2018;2018:4202091.
 - 50) Vizzutti F, Arena U, Romanelli RG, Rega L, Foschi M, Colagrande S, et al. Liver stiffness measurement predicts severe portal hypertension in patients with HCV-related cirrhosis. *Hepatology* 2007;45:1290-1297.
 - 51) Castera L, Pinzani M, Bosch J. Non invasive evaluation of portal hypertension using transient elastography. *J Hepatol* 2012;56:696-703.
 - 52) Augustin S, Millán L, González A, Martell M, Gelabert A, Segarra A, et al. Detection of early portal hypertension with routine data and liver stiffness in patients with asymptomatic liver disease: a prospective study. *J Hepatol* 2014;60:561-569.
 - 53) European Association for the Study of the Liver; Clinical Practice Guideline Panel; Chair; EASL Governing Board representative; Panel members. EASL Clinical Practice Guidelines on non-invasive tests for evaluation of liver disease severity and prognosis – 2021 update. *J Hepatol* 2021;75:659-689.
 - 54) Shi K-Q, Fan Y-C, Pan Z-Z, Lin X-F, Liu W-Y, Chen Y-P, et al. Transient elastography: a meta-analysis of diagnostic accuracy in evaluation of portal hypertension in chronic liver disease. *Liver Int* 2013;33:62-71.
 - 55) You M-W, Kim KW, Pyo J, Huh J, Kim HJ, Lee SJ, et al. A meta-analysis for the diagnostic performance of transient elastography for clinically significant portal hypertension. *Ultrasound Med Biol* 2017;43:59-68.
 - 56) Robic MA, Procopet B, Métivier S, Péron JM, Selves J, Vinel JP, et al. Liver stiffness accurately predicts portal hypertension related complications in patients with chronic liver disease: a prospective study. *J Hepatol* 2011;55:1017-1024.
 - 57) Attia D, Schoenemeier B, Rodt T, Negm A, Lenzen H, Lankisch T, et al. Evaluation of liver and spleen stiffness with acoustic radiation force impulse quantification elastography for diagnosing clinically significant portal hypertension. *Ultraschall Med* 2015;36:603-610.
 - 58) Abraldes JG, Bureau C, Stefanescu H, Augustin S, Ney M, Blasco H, et al.; Anticipate Investigators. Noninvasive tools and risk of clinically significant portal hypertension and varices in compensated cirrhosis: the “Anticipate” study. *Hepatology* 2016;64:2173-2184. Erratum in: *Hepatology* 2017;66:304-305.
 - 59) Galizzi HO, Couto CA, Taranto DOL, Araújo SIO, Vilela EG. Accuracy of non-invasive methods/models for predicting esophageal varices in patients with compensated advanced chronic liver disease secondary to nonalcoholic fatty liver disease. *Ann Hepatol* 2021;20:100229.
 - 60) Sousa M, Fernandes S, Proença L, Silva AP, Leite S, Silva J, et al. The Baveno VI criteria for predicting esophageal varices: validation in real life practice. *Rev Esp Enferm Dig* 2017;109:704-707.
 - 61) Moctezuma-Velazquez C, Saffioti F, Tasayco-Huaman S, Casu S, Mason A, Roccarina D, et al. Non-invasive prediction of high-risk varices in patients with primary biliary cholangitis and primary sclerosing cholangitis. *Am J Gastroenterol* 2019;114:446-452.
 - 62) Pons M, Augustin S, Scheiner B, Guillaume M, Rosselli M, Rodrigues SG, et al. Noninvasive diagnosis of portal hypertension in patients with compensated advanced chronic liver disease. *Am J Gastroenterol* 2021;116:723-732.
 - 63) Wong GL, Chan HL, Choi PC, Chan AW, Lo AO, Chim AM, et al. Association between anthropometric parameters and measurements of liver stiffness by transient elastography. *Clin Gastroenterol Hepatol* 2013;11:295-302.e291-293.
 - 64) Wong V-S, Irls M, Wong G-H, Shili S, Chan A-H, Merrouche W, et al. Unified interpretation of liver stiffness measurement by M and XL probes in non-alcoholic fatty liver disease. *Gut* 2019;68:2057-2064.
 - 65) Nedredal GI, Yin M, McKenzie T, Lillegard J, Luebke-Wheeler J, Talwalkar J, et al. Portal hypertension correlates with splenic stiffness as measured with MR elastography. *J Magn Reson Imaging* 2011;34:79-87.
 - 66) Mejias M, Garcia-Pras E, Gallego J, Méndez R, Bosch J, Fernandez M. Relevance of the mTOR signaling pathway in the pathophysiology of splenomegaly in rats with chronic portal hypertension. *J Hepatol* 2010;52:529-539.
 - 67) Bolognesi M, Di Pascoli M, Verardo A, Gatta A. Splanchnic vasodilation and hyperdynamic circulatory syndrome in cirrhosis. *World J Gastroenterol* 2014;20:2555-2563.
 - 68) Buechter M, Manka P, Theysohn JM, Reinboldt M, Canbay A, Kahraman A. Spleen stiffness is positively correlated with HVPG and decreases significantly after TIPS implantation. *Dig Liver Dis* 2018;50:54-60.
 - 69) Colecchia A, Colli A, Casazza G, Mandolesi D, Schiumerini R, Reggiani LB, et al. Spleen stiffness measurement can predict clinical complications in compensated HCV-related cirrhosis: a prospective study. *J Hepatol* 2014;60:1158-1164.
 - 70) Song J, Huang J, Huang H, Liu S, Luo Y. Performance of spleen stiffness measurement in prediction of clinical significant portal hypertension: a meta-analysis. *Clin Res Hepatol Gastroenterol* 2018;42:216-226.
 - 71) Stefanescu H, Rusu C, Lupsor-Platon M, Nicoara Farcau O, Fischer P, Grigoras C, et al. Liver stiffness assessed by ultrasound shear wave elastography from General Electric accurately predicts clinically significant portal hypertension in patients with advanced chronic liver disease. *Ultraschall Med* 2020;41:526-533.
 - 72) Colecchia A, Ravaoli F, Marasco G, Colli A, Dajti E, Di Biase AR, et al. A combined model based on spleen stiffness

- measurement and Baveno VI criteria to rule out high-risk varices in advanced chronic liver disease. *J Hepatol* 2018;69:308-317.
- 73) Dajti E, Marasco G, Ravaioli F, Alemanni LV, Rossini B, Colecchia L, et al. The role of liver and spleen elastography in advanced chronic liver disease. *Minerva Gastroenterol (Torino)* 2021;67:151-163.
 - 74) Talwalkar JA, Yin M, Venkatesh S, Rossman PJ, Grimm RC, Manduca A, et al. Feasibility of in vivo MR elastographic splenic stiffness measurements in the assessment of portal hypertension. *AJR Am J Roentgenol* 2009;193:122-127.
 - 75) Fierbinteanu-Braticevici C, Tribus L, Peagu R, Petrisor A, Baicus C, Cretoiu D, et al. Spleen stiffness as predictor of esophageal varices in cirrhosis of different etiologies. *Sci Rep* 2019;9:16190.
 - 76) Kim HY, So YH, Kim W, Ahn D-W, Jung YJ, Woo H, et al. Non-invasive response prediction in prophylactic carvedilol therapy for cirrhotic patients with esophageal varices. *J Hepatol* 2019;70:412-422.
 - 77) Ma X, Wang LE, Wu H, Feng Y, Han X, Bu H, et al. Spleen stiffness is superior to liver stiffness for predicting esophageal varices in chronic liver disease: a meta-analysis. *PLoS One* 2016;11:e0165786.
 - 78) Rosołowski M, Hartleb M, Marek T, Milewski J, Linke K, Wallner G, et al. Therapeutic and prophylactic management of bleeding from oesophageal and gastric varices - recommendations of the Working Group of the National Consultant for Gastroenterology. *Prz Gastroenterol* 2014;9:63-68.
 - 79) Jansen C, Möller P, Meyer C, Kolbe CC, Bogs C, Pohlmann A, et al. Increase in liver stiffness after transjugular intrahepatic portosystemic shunt is associated with inflammation and predicts mortality. *Hepatology* 2018;67:1472-1484.
 - 80) Königshofer P. Distinct structural and dynamic components of portal hypertension in different animal models and human liver disease etiologies. *Hepatology* 2021. <https://doi.org/10.1002/hep.32220>.
 - 81) Marasco G, Dajti E, Ravaioli F, Alemanni LV, Capuano F, Gjini K, et al. Spleen stiffness measurement for assessing the response to β -blockers therapy for high-risk esophageal varices patients. *Hepatol Int* 2020;14:850-857.
 - 82) Kim HP, Moon AM, Dellon ES. Noninvasively predicting hemodynamic response to carvedilol in cirrhotic patients with varices: you have some ex-spleening to do. *Gastroenterology* 2019;156:2347-2349.
 - 83) Salzl P, Reiberger T, Ferlitsch M, Payer B, Schwengerer B, Trauner M, et al. Evaluation of portal hypertension and varices by acoustic radiation force impulse imaging of the liver compared to transient elastography and AST to platelet ratio index. *Ultraschall Med* 2014;35:528-533.
 - 84) Elkrief L, Rautou P-E, Ronot M, Lambert S, Dioguardi Burgio M, Francoz C, et al. Prospective comparison of spleen and liver stiffness by using shear-wave and transient elastography for detection of portal hypertension in cirrhosis. *Radiology* 2015;275:589-598.
 - 85) Thiele M, Hugger MB, Kim Y, Rautou P-E, Elkrief L, Jansen C, et al. 2D shear wave liver elastography by Aixplorer to detect portal hypertension in cirrhosis: an individual patient data meta-analysis. *Liver Int* 2020;40:1435-1446.
 - 86) Procopet B, Berzigotti A, Abralde JG, Turon F, Hernandez-Gea V, García-Pagán JC, et al. Real-time shear-wave elastography: applicability, reliability and accuracy for clinically significant portal hypertension. *J Hepatol* 2015;62:1068-1075.
 - 87) Trebicka J, Gu W, de Ledinghen V, Aubé C, Krag A, Praktiknjo M, et al. Two-dimensional shear wave elastography predicts survival in advanced chronic liver disease. *Gut* 2021; <https://doi.org/10.1136/gutjnl-2020-323419>.
 - 88) Wong GLH, Kwok R, Hui AJ, Tse Y-K, Ho K-T, Lo AOS, et al. A new screening strategy for varices by liver and spleen stiffness measurement (LSSM) in cirrhotic patients: a randomized trial. *Liver Int* 2018;38:636-644.
 - 89) Wong G-H, Liang LY, Kwok R, Hui AJ, Tse Y-K, Chan H-Y, et al. Low risk of variceal bleeding in patients with cirrhosis after variceal screening stratified by liver/spleen stiffness. *Hepatology* 2019;70:971-981.
 - 90) Jansen C, Bogs C, Verlinden W, Thiele M, Möller P, Görtzen J, et al. Algorithm to rule out clinically significant portal hypertension combining shear-wave elastography of liver and spleen: a prospective multicentre study. *Gut* 2016;65:1057-1058.
 - 91) Ferraioli G, Tinelli C, Lissandrin R, Zicchetti M, Bernuzzi S, Salvaneschi L, et al. Ultrasound point shear wave elastography assessment of liver and spleen stiffness: effect of training on repeatability of measurements. *Eur Radiol* 2014;24:1283-1289.
 - 92) Castéra L, Foucher J, Bernard P-H, Carvalho F, Allaix D, Merrouche W, et al. Pitfalls of liver stiffness measurement: a 5-year prospective study of 13,369 examinations. *Hepatology* 2010;51:828-835.
 - 93) Ferraioli G, Barr RG. Ultrasound liver elastography beyond liver fibrosis assessment. *World J Gastroenterol* 2020;26:3413-3420.
 - 94) DiPaola FW, Schumacher KR, Goldberg CS, Friedland-Little J, Parameswaran A, Dillman JR. Effect of Fontan operation on liver stiffness in children with single ventricle physiology. *Eur Radiol* 2017;27:2434-2442.
 - 95) Dietrich C, Bamber J, Berzigotti A, Bota S, Cantisani V, Castera L, et al. EFSUMB guidelines and recommendations on the clinical use of liver ultrasound elastography, update 2017 (short version). *Ultraschall Med* 2017;38:377-394.
 - 96) Millonig G, Reimann FM, Friedrich S, Fonouni H, Mehrabi A, Büchler MW, et al. Extrahepatic cholestasis increases liver stiffness (FibroScan) irrespective of fibrosis. *Hepatology* 2008;48:1718-1723.
 - 97) Berzigotti A, De Gottardi A, Vukotic R, Siramolpiwat S, Abralde JG, García-Pagan JC, et al. Effect of meal ingestion on liver stiffness in patients with cirrhosis and portal hypertension. *PLoS One* 2013;8:e58742.
 - 98) Alvarez D, Orozco F, Mella JM, Anders M, Antinucci F, Mastai R. Meal ingestion markedly increases liver stiffness suggesting the need for liver stiffness determination in fasting conditions. *Gastroenterol Hepatol* 2015;38:431-435.
 - 99) Kjærgaard M, Thiele M, Jansen C, Stæhr Madsen B, Görtzen J, Strassburg C, et al. High risk of misinterpreting liver and spleen stiffness using 2D shear-wave and transient elastography after a moderate or high calorie meal. *PLoS One* 2017;12:e0173992.
 - 100) Yoo J-J, Seo YS, Kim YS, Jeong SW, Jang JY, Suh SJ, et al. The influence of histologic inflammation on the improvement of liver stiffness values over 1 and 3 years. *J Clin Med* 2019;8:2065.
 - 101) Mueller S, Englert S, Seitz HK, Badea RI, Erhardt A, Bozaari B, et al. Inflammation-adapted liver stiffness values for improved fibrosis staging in patients with hepatitis C virus and alcoholic liver disease. *Liver Int* 2015;35:2514-2521.
 - 102) Janik MK, Kruk B, Szczepankiewicz B, Kostrzewa K, Raszeja-Wyszomirska J, Górnicka B, et al. Measurement of liver and spleen stiffness as complementary methods for assessment of liver fibrosis in autoimmune hepatitis. *Liver Int* 2021;41:348-356.
 - 103) Giuffrè M, Fouraki S, Campigotto M, Colombo A, Visintin A, Buonocore MR, et al. Alanine aminotransferase and splenoportal dynamics affect spleen stiffness measured by point shear-wave elastography in patients with chronic hepatitis C in the absence of significant liver fibrosis. *J Ultrasound* 2021;24:67-73.
 - 104) Wang H, Wen B, Chang X, Wu Q, Wen W, Zhou F, et al. Baveno VI criteria and spleen stiffness measurement rule out high-risk varices in virally suppressed HBV-related cirrhosis. *J Hepatol* 2021;74:584-592.
 - 105) Bureau C, Metivier S, Peron JM, Selves J, Robic MA, Gourraud PA, et al. Transient elastography accurately predicts presence of

- significant portal hypertension in patients with chronic liver disease. *Aliment Pharmacol Ther* 2008;27:1261-1268.
- 106) Lemoine M, Katsahian S, Ziol M, Nahon P, Ganne-carrie N, Kazemi F, et al. Liver stiffness measurement as a predictive tool of clinically significant portal hypertension in patients with compensated hepatitis C virus or alcohol-related cirrhosis. *Aliment Pharmacol Ther* 2008;28:1102-1110.
- 107) Sánchez-Conde M, Montes-Ramírez ML, Miralles P, Castro Alvarez JM, Bellón JM, Ramírez M, et al. Comparison of transient elastography and liver biopsy for the assessment of liver fibrosis in HIV/hepatitis C virus-coinfected patients and correlation with noninvasive serum markers. *J Viral Hepat* 2010;17:280-286.
- 108) Reiberger T, Ferlitsch A, Payer BA, Pinter M, Schwabl P, Stift J, et al. Noninvasive screening for liver fibrosis and portal hypertension by transient elastography—a large single center experience. *Wien Klin Wochenschr* 2012;124:395-402.
- 109) Llop E, Berzigotti A, Reig M, Erice E, Reverter E, Seojo S, et al. Assessment of portal hypertension by transient elastography in patients with compensated cirrhosis and potentially resectable liver tumors. *J Hepatol* 2012;56:103-108.
- 110) Hong WK, Kim MY, Baik SK, Shin SY, Kim JM, Kang YS, et al. The usefulness of non-invasive liver stiffness measurements in predicting clinically significant portal hypertension in cirrhotic patients: Korean data. *Clin Mol Hepatol* 2013;19:370-375.
- 111) Sharma P, Kirnake V, Tyagi P, Bansal N, Singla V, Kumar A, et al. Spleen stiffness in patients with cirrhosis in predicting esophageal varices. *Am J Gastroenterol* 2013;108:1101-1107.
- 112) Kim TY, Jeong WK, Sohn JH, Kim J, Kim MY, Kim Y. Evaluation of portal hypertension by real-time shear wave elastography in cirrhotic patients. *Liver Int* 2015;35:2416-2424.
- 113) Schwabl P, Bota S, Salzl P, Mandorfer M, Payer BA, Ferlitsch A, et al. New reliability criteria for transient elastography increase the number of accurate measurements for screening of cirrhosis and portal hypertension. *Liver Int* 2015;35:381-390.
- 114) Jansen C, Bogs C, Verlinden W, Thiele M, Möller P, Görtzen J, et al. Shear-wave elastography of the liver and spleen identifies clinically significant portal hypertension: a prospective multicentre study. *Liver Int* 2017;37:396-405.
- 115) Souhami A, Sartoris R, Rautou P-E, Cauchy F, Bouattour M, Durand F, et al. Similar performance of liver stiffness measurement and liver surface nodularity for the detection of portal hypertension in patients with hepatocellular carcinoma. *JHEP Rep* 2020;2:100147.
- 116) Kim HY, Jin EH, Kim W, Lee JY, Woo H, Oh S, et al. The role of spleen stiffness in determining the severity and bleeding risk of esophageal varices in cirrhotic patients. *Medicine (Baltimore)* 2015;94:e1031.
- 117) Marasco G, Colecchia A, Colli A, Ravaioli F, Casazza G, Bacchi Reggiani ML, et al. Role of liver and spleen stiffness in predicting the recurrence of hepatocellular carcinoma after resection. *J Hepatol* 2019;70:440-448.
- 118) Stefanescu H, Marasco G, Calès P, Fraquelli M, Rosselli M, Ganne-Carriè N, et al. A novel spleen-dedicated stiffness measurement by FibroScan® improves the screening of high-risk oesophageal varices. *Liver Int* 2020;40:175-185.
- 119) Cho YS, Kim Y, Sohn JH. Application of supersonic shear imaging to the Baveno VI criteria and a combination model with spleen stiffness measurement to rule out high-risk varices in compensated advanced chronic liver disease. *Ultraschall Med* 2020; <https://doi.org/10.1055/a-1168-6271>.
- 120) Dajti E, Marasco G, Ravaioli F, Colecchia L, Ferrarese A, Festi D, et al. Risk of hepatocellular carcinoma after HCV eradication: determining the role of portal hypertension by measuring spleen stiffness. *JHEP Rep* 2021;3:100289.

Author names in bold designate shared co-first authorship.